



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CALCULUS.

Solutions of Problem 172 have been received from J. E. Sanders, Hackney, Ohio, and J. B. Gregg, M. Sc., C. E., Senecaville, Ohio.

173. Proposed by J. E. SANDERS, Hackney, Ohio.

Find the greatest ellipse that can be inscribed in a quadrant of a given circle.

Solution by J. B. GREGG, M. Sc., C. E., Senecaville, Ohio, and G. W. GREENWOOD, B. A. (Oxon), Professor of Mathematics and Astronomy, McKendree College, Lebanon, Ill.

By symmetry, the radius bisecting the quadrant will coincide with an axis of the ellipse. Denote the length of this semi-axis, and that of the transverse semi-axis, by a , b . Then

$$\pi ab = \text{area of ellipse} = A \text{ (say).}$$

$$a + \sqrt{a^2 + b^2} = \text{radius of given circle} = r \text{ (say).}$$

$$\therefore r^2 = 2ar + b^2.$$

$$\text{Since } A \text{ is a maximum, } bda + adb = 0. \text{ Also } rda + bdb = 0.$$

$$\therefore ra = b^2, r = 3a, A = \frac{1}{3}\pi r^2 \sqrt{3}.$$

Also solved by G. B. M. Zerr, A. M., Ph. D., Parsons, W. Va., and A. H. Holmes, Brunswick, Me.

MECHANICS.

163. Proposed by W. J. GREENSTREET, A. M., Editor of The Mathematical Gazette, Stroud, England.

A particle A , mass m , rests on a smooth horizontal plane and is attached by two inelastic strings to masses m_1 , m_2 at points B and C such that BAC is a right angle. If a blow is given A at an angle θ to AB , find the initial direction of motion of m , and equations for initial motion of the particles m_1 and m_2 .

Solution by J. B. GREGG, M. Sc., C. E., Senecaville, Ohio.

Let v , v_1 , and v_2 be the respective initial velocities of m , m_1 , and m_2 , and let ϕ be the angle which the initial motion of m makes with the line of direction of the blow. Construct $AD = v$; then DE perpendicular to BA produced $= v_2$, $AE = v_1$, $\angle DAE = \theta + \phi$.

$$v_1 = v \cos(\theta + \phi), v_2 = v \sin(\theta + \phi), mv \sin \phi + m_1 v_1 \sin \theta = m_2 v_2 \cos \theta.$$

$$\text{Solving for } \phi, \tan \phi = \frac{(m_2 - m_1) \sin \theta \cos \theta}{m - m_1 \sin^2 \theta - m_2 \cos^2 \theta}.$$

DIOPHANTINE ANALYSIS.

116. Proposed by HARRY S. VANDIVER, Bala, Pa.

If n is an odd positive integer, and $1, n, n', n'', \dots$ denote all its distinct divisors, then $2^n > 2(n+1)(n'+1)(n''+1)\dots$

II. Solution by the PROPOSER.

By Euler's generalization of Fermat's Theorem (Dirichlet-Dedekind Zah-